arXiv:quant-ph/0501179v1 31 Jan 2005

Comments on "Evidence for Quantized Displacement in Macroscopic Nanomechanical Oscillators"

A. Kwang-Hua CHU

P.O. Box 30-15, Shanghai 200030, PR China

and

Department of Physics,

Xinjiang University, 14, Road Shengli,

Urumqi 830046, PR China

Abstract

We make comments on Gaidarzhy *et al.*'s [*Phys. Rev. Lett.* 94, 030402 (2005)] letter.

PACS numbers: 03.65.Ta, 62.25.+g, 62.30.+d, 62.40.+i

Keywords:

Quite recently Gaidarzhy *et al.* [1] have measured displacement in a series of nanomechanical oscillators [2-3] with 1.49-GHz resonance frequencies down to 110 mK temperatures. While the low frequency mode at 21 MHz shows classical behavior with expected drive dependence ($\propto B^2$), the 1.49 GHz mode displays nonmonotonic dependence on driving force at a temperature which corresponds to a thermal occupation number $N_{th} \rightarrow 1$ (cf. Fig. 4 therein [1]). They finally claimed that their experimental data indicate the first observation of quantum displacement in macroscopic nanomechanical oscillators. However, as they also argued that, there may be other semiclassical mechanisms which involve both nonlinearity and quantum effects [4].

The quantum mechanical harmonic oscillator [1-3] is a fundamental example of textbook quantum mechanics. Understanding the measurement of the quantum system itself requires identifying the components of the quantum measurement process: quantum system, measuring apparatus, and their interaction [5]. Coupling to the environment, which introduces decoherence and dissipation, must be included in the theoretical analysis [6].

Based on the present author's experiences, we like to make some remarks about the possible origin of their so-called *quantum displacement* in macroscopic nanomechanical oscillators. Firstly, judged from the brief introduction of their experimental procedure, their might be no explicit Casmir effects [7]. However, the zero-point vibrations at low temperature and the anharmonic decay of the phonon [8] should be carefully taken into account as the authors already mentioned : the intermediate jumps between these two (discrete) states (in the range of $6.5 - 7$ Tesla in Fig. 4(b) [1]) could represent transitions induced by thermal fluctuations, as the temperature even at $N_{th} \rightarrow 1$ is high enough that the thermal energy k_BT smears the gap hf in the energy levels.

On the other hand, as noted by Gaidarzhy *et al.* in [1] : another possibility is that the oscillator does not start from the energy eigenbasis, as there is no *a priori* reason for it to be in this preferred basis. The present author would like to point out, the stochastic resonance [9-10], e.g., the two-state model [10], might also induce the results reported in Fig. 4 of [1]. Either the stochastic resonance in a paramagnetically driven bistable buckling ribbon [11] or a magnetostochastic resonance in thin films [12] have been reported previously. To be specific, considering Fig. 4 (c) in [1], as mentioned therein : two sweeps of the magnetic field, up (black) and down (blue), reproducing the same transition, with occasional transitions to a possible intermediate state, there also are possible weak stochastic resonances in the range of $10 - 10.5$ Tesla (cf. Fig. 4 (c) in [1] together with Figs. 3 and 4 of Gammaitoni *et al.* in [10]).

In brief, detailed considerations of Casmir effects together with anharmonicity of thermal phonons as well as stochastic resonances should be made to draw the final conclusion for the possible observation of quantum displacement in [1].

---

[1] A. Gaidarzhy, G. Zolfagharkhani, R.L. Badzey, and P. Mohanty, Phys. Rev. Lett. 94, 030402 (2005).

[2] M. Blencowe, Phys. Rep. 395, 159 (2004).

[3] R.H. Blick, F.W. Beil, E. Höhberger, A. Erbe, C. Weiss, Nano-Electromechanical Systems: Displacement Detection and the Mechanical Single Electron Shuttle, in *Lecture Notes in Physics* (vol. 579, Springer, Berlin, 2001) pp. 215.

[4] V. Peano and M. Thorwart, Phys. Rev. B 70, 235401 (2004).

[5] W. Zurek, Rev. Mod. Phys. 75, 715 (2003).

[6] A. J. Leggett, J. Phys. Condens. Matter 14, R415 (2002). A. J. Leggett, Phys. Scr. T102, 69 (2002). P. A. M. Dirac, The Principles of Quantum Mechanics (Clarendon , Oxford, 1981).

[7] K.A. Milton, J. Phys. A : Math. Gen. 37, R209-R277 (2004).

[8] M. Cardona, Ann. Phys. (Leipzig) 9, 865 (2000).

[9] T. Wellens, V. Shatokhin, and A. Buchleitner, Rep. Prog. Phys. 67, 45-105 (2004).

[10] L. Gammaitoni, P. Hänggi, P. Jung and F. Marchesoni, Rev. Mod. Phys. 70, 223 (1998). Z. K.-H. Chu, Eur. J. Phys. 21, L19-L20 (2000). *ibid* 23, L23-L24 (2002).

[11] M.L. Spano, M. Wun-Fogle, and W.L. Ditto, Phys. Rev. A 46, R5253 (1992).

[12] A.N. Grigorenko, P.I. Nikitin, A.N. Slavin, and P.Y. Zhou, J. Appl. Phys. 76, 6335 (1994).